



Year: 2018

The rationale and design of the SMART CRT trial

Gold, Michael R ; Auricchio, Angelo ; Leclercq, Christophe ; Lowy, Jonathan ; Rials, Seth J ; Shoda, Morio ; Tomassoni, Gery ; Yong, Patrick ; Wold, Nicholas ; Ellenbogen, Kenneth A

Abstract: AIMS The SMART CRT study will assess the efficacy of an atrioventricular optimization algorithm to improve reverse remodeling among patients undergoing cardiac resynchronization therapy (CRT) in the presence of interventricular electrical delay. **METHODS AND RESULTS** The SMART CRT study is a global, multicenter, prospective, randomized study of patients undergoing CRT implantation. The primary endpoint of this trial is response rate to CRT, defined as decrease in left ventricular end-systolic volume (LVESV) 15% at 6 months compared to preimplant baseline. Additional prespecified analyses are: (1) clinical composite endpoint combining all-cause mortality, heart failure events, New York Heart Association class, and Quality of Life (using a patient global assessment instrument); (2) the individual components of the clinical composite endpoint; (3) 6-minute walk distance; (4) Kansas City Cardiomyopathy Questionnaire; (5) LVESV as a continuous variable; and (6) absolute left-ventricular ejection fraction. Subjects with intraventricular delay 70 ms measured between the right ventricular and left ventricular pacing leads will be randomized in a 1:1 ratio to have either an AV Delay and pacing chamber determined by SmartDelay™ or a Fixed AV Delay of 120 ms with biventricular pacing. Enrollment of an estimated 726 of subjects from up to 100 centers worldwide is planned to achieve 436 randomized subjects and 370 complete data sets required to power the primary endpoint. **CONCLUSIONS** This trial will provide important data regarding the importance of AV Delay programming in patients with prolonged interventricular delay at the pacing sites.

DOI: <https://doi.org/10.1111/pace.13459>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-158724>

Journal Article

Accepted Version

Originally published at:

Gold, Michael R; Auricchio, Angelo; Leclercq, Christophe; Lowy, Jonathan; Rials, Seth J; Shoda, Morio; Tomassoni, Gery; Yong, Patrick; Wold, Nicholas; Ellenbogen, Kenneth A (2018). The rationale and design of the SMART CRT trial. *Pacing and Clinical Electrophysiology : PACE*, 41(9):1212-1216.

DOI: <https://doi.org/10.1111/pace.13459>

The Rationale and Design of the SMART CRT Trial

Short Title: SMART CRT Study Design

Michael R. Gold, PhD, MD¹, Angelo Auricchio, MD, PhD², , Christophe Leclercq, MD, PhD³,
Jonathan Lowy, MD⁴, Seth Rials, MD, PhD⁵, Morio Shoda, MD, PhD⁶, Gery Tomassoni, MD⁷,
Patrick Yong⁸, Nicholas Wold⁸, Kenneth A. Ellenbogen, MD⁹

¹Medical University of South Carolina, Charleston, SC, USA, ²Cardiocentro Ticino, Lugano, Switzerland, , ³CHRU Hopital Pontchaillou, France, ⁴PeaceHealth Southwest Medical, Vancouver, WA, USA, ⁵OhioHealth Research and Innovation Institute, Columbus, OH, USA, ⁶Tokyo Medical University Hospital, Tokyo, Japan ⁷Baptist Health Lexington, Lexington, KY, USA ⁸Boston Scientific Corp., St Paul, MN, USA, ⁹Virginia Commonwealth University, Richmond, VA, USA

Position, Institution & location of all authors:

Dr. Gold: Micheal E Assey Professor, Department of Medicine, Medical University of South Carolina, SC, USA.

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/pace.13459](#).

This article is protected by copyright. All rights reserved.

Dr. Auricchio: Director Clinical Electrophysiology Program, Division of Cardiology, Fondazione
Cardiocentro Ticino, Lugano, Switzerland

Dr. Leclercq: CHRU Hopital Pontchaillou, France

Dr. Lowy: Medical Director of Cardiac Research, PeaceHealth Southwest Medical Center,
Vancouver, WA, USA

Dr. Rials: OhioHealth Research and Innovation Institute, Columbus, OH, USA

Dr. Shoda: Tokyo Medical University Hospital, Tokyo, Japan

Dr. Tomassoni: Baptist Health Lexington, Lexington, KY, USA

Mr. Yong: Fellow, Clinical Science, Boston Scientific Corp., St. Paul, MN

Mr. Wold: Principal Biostatistician, Clinical Science, Boston Scientific Corp., St. Paul, MN

Dr. Ellenbogen: Martha and Harold Kimmerling Professor, Department of Medicine, Virginia
Commonwealth University School of Medicine, Richmond, VA

Corresponding Author:

Dr. Michael R. Gold

Mailing Address: 30 Courtenay Drive,, MSC 592, Charleston, SC 29425 USA

Telephone: (843) 876-4760

Fax: (843) 876-4990

Email: goldmr@musc.edu

Abstract:

Aims: The SMART CRT study will assess the efficacy of an AV optimization algorithm to improve reverse remodeling among patients undergoing cardiac resynchronization therapy in the presence of interventricular electrical delay.

Methods and Results: The SMART CRT study is a global, multi-center, prospective, randomized study of patients undergoing cardiac resynchronization therapy (CRT) implantation. The primary endpoint of this trial is response rate to CRT, defined as decrease in left ventricular end systolic volume (LVESV) $\geq 15\%$ at 6 months compared to pre-implant baseline. Additional prespecified analyses are: a) clinical composite endpoint combining all-cause mortality, heart failure events, NYHA class and Quality of Life (using a patient global assessment instrument), b) the individual components of the clinical composite endpoint, c) 6-minute walk distance, d) Kansas City Cardiomyopathy Questionnaire e) LVESV as a continuous variable, and f) absolute left-ventricular ejection fraction (LVEF). Subjects with intraventricular delay ≥ 70 msec measured between the right ventricular and left ventricular pacing leads will be randomized in a 1:1 ratio to have either an AV Delay and pacing chamber determined by SmartDelay™ or a Fixed AV Delay of 120 msec with biventricular pacing. Enrollment of an estimated 726 of subjects from up to 100 centers worldwide is planned to achieve 436 randomized subjects and 370 complete datasets required to power the primary endpoint.

Conclusions: This trial will provide important data regarding the importance of AV Delay programming in patients with prolonged interventricular delay at the pacing sites.

ClinicalTrials.gov Identifier: NCT03089281

Key words: Heart Failure; Cardiac Resynchronization Therapy; Clinical Trials;

Abbreviations:

AVO – Atrio-ventricular optimization

BSC – Boston Scientific Corporation

CRT – Cardiac resynchronization therapy

ECG – Electrocardiogram

HF – Heart failure

KCCQ – Kansas City Cardiomyopathy Questionnaire

LV – Left ventricular

LVEF – Left ventricular ejection fraction

LVESV – Left ventricular end systolic volume

NYHA – New York Heart Association classification.

Introduction

Cardiac resynchronization therapy (CRT) is a well-established treatment for patients with symptomatic heart failure (HF), a reduced left ventricular (LV) ejection fraction (LVEF) and QRS prolongation. CRT has a number of beneficial effects including improvement in measures of quality of life, reverse LV remodeling and reductions in HF hospitalizations and mortality.¹⁻⁵ The magnitude of benefit with CRT is affected by patient characteristics such as QRS duration and morphology, etiology of HF and gender. More recently it has been shown that procedural factors, such as LV pacing at sites of late electrical activation, as assessed by either the QLV or RV-LV durations, are associated with improved CRT response.⁶⁻¹¹

Optimizing atrio-ventricular (AV) timing was a standard part of CRT management for many years, but routine AV optimization (AVO) is no longer recommended based on disappointing results from multicenter randomized trials.¹²⁻¹⁴ However, retrospective analyses have shown that certain subgroups of patients may benefit from AVO, including those paced at sites of increased electrical delay.¹⁵ To understand better the role of AVO among such patients, the SMART CRT study was designed as a prospective, randomized trial among patients paced with prolonged interventricular delay.

Rationale

A post hoc analysis of the SMART-AV trial showed that increasing RV-LV duration was associated with greater reverse remodeling with AVO compared with nominal programming.¹¹ For the analysis, the patient population was assessed by grouping into quartiles based on RV-LV duration. To provide more granularity to the data for the design of SMART CRT the splitting variable was changed to 10 msec increments. As shown in Panel A of Figure 1, the proportion of patients with an RV-LV duration greater than a given cutoff value diminishes as the RV-LV duration increases (median duration was 70 msec). The patient population was evaluated in a similar fashion in which the

response rates for AVO and nominal programming were compared. As the RV-LV duration increases, the difference in response rates also increases (Panel B of Figure 1). The curve is nonlinear with a plateau area. Choosing a large cutoff value increases the effect size but also diminishes the number of available patients. To maximize the proportion of patients who may benefit from AVO, enrolling patients with an RV-LV duration near the onset of the plateau region (70 msec) was chosen. This value ensures that a clinically meaningful improvement in response rate would be observed while ensuring that the cutoff value is applicable to the majority of patients.

Study Hypotheses

The primary hypothesis of SMART CRT is that among patients with prolonged inrerventricular delay between the RV and LV leads, CRT with AVO will result in greater reverse LV remodeling compared with CRT programmed at nominal settings.

Study Design

The SMART CRT study is a global, multi-center, prospective, randomized study that will enroll subjects who intend to undergo *de novo* implantation of a CRT System (Boston Scientific, Marlborough, MA) or upgrade from a preexisting single or dual chamber device. Only patients with at least one RV-LV duration ≥ 70 msec between programmable RV lead and LV pacing configuration, as measured automatically from the pulse generator, will be randomized. An acceptable pacing configuration is one with a pacing threshold ≤ 4.5 V at a pulse width of 0.5 msec and no diaphragmatic stimulation within 3 V above the pacing threshold. If multiple vectors meet the acceptable pacing criteria, the investigator will select a vector with an RV-LV delay within 10 msec

of the longest measured delay. In cases where there is no acceptable pacing configuration that meets randomization, the right or left ventricular lead can be repositioned at the discretion of the implanting physician; otherwise, the patient will be exited from the protocol. Subjects will be randomized in a 1:1 ratio to have either an AV Delay and pacing chamber determined by SmartDelay™ or a Fixed AV Delay of 120 msec with biventricular pacing. Of note, multisite pacing is not allowed in the protocol. Randomized subjects will be followed semi-annually for a minimum of 540 days \pm 60 days window (i.e., 18 months) post implant. Figure 2 is a flow chart which describes the proposed study plan from first enrollment to study closure. All subjects will receive a *de novo* quadripolar Boston Scientific Cardiac Resynchronization Therapy Defibrillator (CRT-D) in conjunction with an AUCITY X4 LV lead. Lead locations will be at the discretion of the implanting physician.

Study Population

The study population consists of patients with a CRT indication scheduled for either *de novo* implantation or upgrade from single or dual chamber device to a CRT-D. Of note, given that multiple geographies were included in this trial it was decided to use as uniform as possible an inclusion criteria rather than individual guidelines of the various regions.

The inclusion and exclusion criteria are further described in detail within Tables 1 and 2, respectively.

Study Endpoints

The primary endpoint of this trial is proportion of patients with a positive response to CRT. CRT response is defined as a decrease in left ventricular end systolic volume (LVESV) \geq 15% at 6 months compared to pre-implant baseline. The proportion of patients responding to CRT will be compared between the randomized groups. In addition to the formal endpoint hypotheses, comparisons between groups for pre-specified analyses include, but are not limited to: a) clinical

composite endpoint combining all-cause mortality, heart failure events, NYHA class and Quality of Life (using a patient global assessment instrument), b) the individual components of the clinical composite endpoint, c) 6-minute walk distance, d) Kansas City Cardiomyopathy Questionnaire (KCCQ), e) LVESV as a continuous variable, and f) LVEF.

Trial Oversight

The study protocol was developed with a steering committee consisting of 8 physicians and a patient advocate. The local ethics committees of all participating sites are required to approve the study protocol. Central and onsite monitoring, data review and safety oversight will be provided by the sponsor, Boston Scientific (Marlborough, MA, USA). Echocardiograms will be interpreted by a blinded core laboratory. All heart failure events and deaths will be reviewed by an independent clinical event committee.

Power Calculations

For the primary endpoint difference in response rates between the group with a fixed AV Delay (120ms) and AV Delay as specified by SmartDelay the sample size is powered to obtain a 90% confidence level with a one-sided alpha of 2.5%. It is expected that 60% of enrolled patients will have an RV-LV-interval of ≥ 70 msec and undergo randomization. Attrition after randomization is expected to be 15%. Under those assumptions, an estimated 726 subjects will be enrolled to achieve 436 randomized subjects and 370 patients in the analysis cohort.

Statistical Analyses

The Primary Endpoint, in addition to the additional analyses, will be evaluated using intention-to-treat methodology, in which subjects with complete datasets are evaluated according to their randomized treatment groups. The Primary Endpoint will be evaluated using a chi-square test with a one-sided significance level of 2.5%, comparing the proportion of responders in the SmartDelay randomized group to the Fixed AV Delay randomized group. The following sensitivity analyses for the Primary Endpoint will be performed: intention-to-treat with missing endpoint data imputed via multiple imputation, on-treatment in which subjects are analyzed according treatment received and per protocol including only subjects programmed to their randomized assignment are included.

The clinical composite endpoint will be assessed utilizing a Cochran-Armitage test for trend. Cochran–Mantel–Haenszel tests will be used to evaluate NYHA classification and patient global assessment. The continuous outcomes of 6-minute walk distance, KCCQ, LVESV and LVEF will be tested using a two-sample t-test. The time-to-event analyses of all-cause mortality and heart failure events will be evaluated using a log-rank test; data from any subjects who are event-free will be right censored on the final date of study participation.

Subgroup analyses for the Primary Endpoint will be performed by testing the interaction between the subgroup and randomized group utilizing a logistic regression model. Subgroup analyses will be performed on the following characteristics: age, gender, ischemic etiology, conduction disorder, QRS duration, LV lead location, baseline NYHA classification and SmartDelay recommended programming (LV only vs. biventricular pacing, and AV Delay equal to 120 ± 20 msec vs. all others). Multivariable logistic regression modeling will also be performed using these characteristics.

An interim sample size re-estimation will be performed after at least half of the required 6 month datasets ($N = 185$) have been completed and reviewed by the core lab. The pooled and blinded response rate across both groups will be assessed in this interim sample size re-assessment cohort.

The observed rate will be weighted by the number of patients in each group and compared to the assumed weighted rate. If the observed rate is less than the assumed rate, the number of patients required for randomization will be increased accordingly, up to a maximum of 456 randomized subjects. The treatment effect will not be calculated, thereby maintaining the blind.

Discussion

Despite early enthusiasm for AV optimization, primarily using echocardiographic techniques, current consensus documents and guidelines do not recommend routine optimization based on the results of prospective, multicenter trials.^{16,17} As such, routine AVO has fallen out of favor. However, analyses of the SMART-AV trial showed that the electrogram based algorithm (SmartDelay) was more effective among patients with the LV lead at sites with a long left ventricular (i.e. QLV)¹⁵ or interventricular (RV-LV) delay.¹¹ The present trial is designed to confirm these findings prospectively by only randomizing patients with a prolonged RV-LV duration at implant. This would support the hypothesis that AVO is most effective among patients more likely to respond to CRT rather than the initial hope of AVO to “convert” non-responders to responders.

The Adaptive CRT trial was another randomized trial which showed no benefit of AVO for the whole CRT cohort,¹³ but a subgroup was identified with LBBB and normal PR interval that was associated with an improved response.¹⁸ The AdaptResponse trial is a prospective study designed to validate this finding.¹⁹ There are some similarities between these algorithms as both are designed to promote fusion of intrinsic conduction to produce more synchronous LV activation including LV only pacing rather than choosing AV delays to optimize LV filling. Since LBBB patients have longer RV-LV times,¹⁰ SMART CRT and AdaptResponse may evaluate a common mechanistic hypothesis that AVO that promotes electrical resynchronization is of value primarily among patients with electrical delay.

There has been much interest in reducing the purported non-responder rate to CRT. Although there is no consensus on the definition of response to this therapy it is clear that optimizing outcomes is an important goal. In this regard, certain clinical factors such as etiology of HF, gender and ECG characteristics including QRS duration and morphology affect CRT response. More recently, the role of targeting LV pacing to areas of late mechanical or electrical delay has also been shown to impact CRT response. The SMART CRT trial will further add to our understanding of strategies to optimize CRT outcomes if the results show a benefit of combining AVO with pacing at site of prolonged interventricular conduction delay.

In summary, SMART CRT is a prospective, randomized study that will be the first trial to evaluate the SmartDelay algorithm in patients with interventricular conduction delay using a quadripolar left ventricular transvenous lead. At present there are no multicenter, randomized, blinded trials showing the benefit of AVO for CRT so this study will potentially establish the benefit of routine AVO in a subset of CRT patients.

Word Count (manuscript text, not including abstract, acknowledgements and disclosures. Maximum 2500 words, 20 references, 5 figures or tables): **2120**

Acknowledgements

Funding

The SMART CRT study is supported and funded by Boston Scientific Corporation.

Institutional Affiliations of Authors (Conflict of Interest):

Dr. Gold: research grants and consulting from Boston Scientific

Dr. Auricchio: consulting for Boston Scientific; consulting for Liva Nova

Dr. Rials: honorarium from Boston Scientific

Dr. Shoda: research grants from Abbott; research grants from Biotronik; research grants from Boston Scientific; research grants from Medtronic

Dr. Tomassoni: consulting and honorarium from Abbott; consulting and honorarium from Biotronik; consulting and honorarium from Boston Scientific; consulting and honorarium from Medtronic

Mr. Yong and Mr. Wold: employee of Boston Scientific

Dr. Ellenbogen: consulting and honorarium from Abbott; honorarium from Biotronik; consulting, honorarium, and research grants from Boston Scientific; consulting, honorarium, and research grants from Medtronic.

Dr. Lowy, Dr. Leclercq: none

Author Contributions:

Dr. Gold: concept/study design, critical revision and drafting the article

Dr. Auricchio: concept/study design, critical revision of the article

Dr. Leclercq: concept/study design, critical revision of the article

Dr. Lowy: concept/study design, critical revision of the article

Dr. Rials: concept/study design, critical revision of the article

Dr. Shoda: concept/study design, critical revision of the article

Dr. Tomassoni: concept/study design, critical revision of the article

Mr. Yong: concept/study design, critical revision of the article

Mr. Wold: concept/study design, critical revision of the article

Dr. Ellenbogen: concept/study design, critical revision of the article

References

1. Bristow MR, Saxon LA, Boehmer J, Krueger S, Kass DA, De Marco T, Carson P, DiCarlo L, DeMets D, White BG, DeVries DW, Feldman AM, Comparison of Medical Therapy, Pacing and D in HF (COMPANION) I. Cardiac-resynchronization therapy with or without an implantable defibrillator in advanced chronic heart failure. *N Engl J Med*. 2004;350:2140–50.
2. Cleland J, Daubert J, Erdmann E, Freemantle N, Gras D, Kappenberger L, Tavazzi L. The effect of cardiac resynchronization on morbidity and mortality in heart failure. *N Engl J Med*. 2005;352:1539–1549.
3. Moss AJ, Hall WJ, Cannom DS, Klein H, Brown MW, Daubert JP, Estes NAM, Foster E, Greenberg H, Higgins SL, Pfeffer MA, Solomon SD, Wilber D, Zareba W, MADIT-CRT Trial Investigators. Cardiac-resynchronization therapy for the prevention of heart-failure events. *N Engl J Med*. 2009;361:1329–38.

4. Abraham WT, Fisher WG, Smith AL, Delurgio DB, Leon AR, Loh E, Kocovic DZ, Packer M, Clavell AL, Hayes DL, Ellestad M, Trupp RJ, Underwood J, Pickering F, Truex C, McAtee P, Messenger J, MIRACLE Study Group. Multicenter InSync Randomized Clinical Evaluation. Cardiac resynchronization in chronic heart failure. *N Engl J Med*. 2002;346:1845–53.
5. Linde C, Abraham WT, Gold MR, St. John Sutton M, Ghio S, Daubert C. Randomized Trial of Cardiac Resynchronization in Mildly Symptomatic Heart Failure Patients and in Asymptomatic Patients With Left Ventricular Dysfunction and Previous Heart Failure Symptoms. *J Am Coll Cardiol*. 2008;52:1834–1843.
6. Singh JP, Fan D, Heist EK, Alabiad CR, Taub C, Reddy V, Mansour M, Picard MH, Ruskin JN, Mela T. Left ventricular lead electrical delay predicts response to cardiac resynchronization therapy. *Heart Rhythm*. 2006;3:1285–1292.
7. Gold MR, Birgersdotter-Green U, Singh JP, Ellenbogen K a., Yu Y, Meyer TE, Seth M, Tchou PJ. The relationship between ventricular electrical delay and left ventricular remodelling with cardiac resynchronization therapy. *Eur Heart J*. 2011;32:2516–2524.
8. D’Onofrio A, Botto G, Mantica M, La Rosa C, Occhetta E, Verlato R, Molon G, Ammendola E, Villani GQ, Bongiorni MG, Gelmini GP, Ciardiello C, Dicandia CD. The interventricular conduction time is associated with response to cardiac resynchronization therapy: Interventricular electrical delay. *Int J Cardiol*. 2013;168:5067–5068.
9. D’Onofrio A, Botto G, Mantica M, La Rosa C, Occhetta E, Verlato R, Molon G, Ammendola E, Villani GQ, Bongiorni MG, Bianchi V, Gelmini GP, Valsecchi S, Ciardiello C. Incremental value of larger interventricular conduction time in improving cardiac resynchronization therapy outcome in patients with different QRS duration. *J Cardiovasc Electrophysiol*.

- 2014;25:500–506.
10. Gold MR, Yu Y, Wold N, Day JD. The role of interventricular conduction delay to predict clinical response with cardiac resynchronization therapy. *Heart Rhythm*. 2017;14:1748–1755.
 11. Gold MR, Singh JP, Ellenbogen KA, Yu Y, Wold N, Meyer TE, Birgersdöter-Green U. Interventricular Electrical Delay Is Predictive of Response to Cardiac Resynchronization Therapy. *JACC Clin Electrophysiol*. 2016;2:438–447.
 12. Ellenbogen KA, Gold MR, Meyer TE, Fernández Lozano I, Mittal S, Waggoner AD, Lemke B, Singh JP, Spinale FG, Van Eyk JE, Whitehill J, Weiner S, Bedi M, Rapkin J, Stein KM. Primary results from the smartdelay determined AV optimization: A comparison to other AV delay methods used in cardiac resynchronization therapy (SMART-AV) trial: A randomized trial comparing empirical, echocardiography- guided, and algorithmic atrioventr. *Circulation*. 2010;122:2660–2668.
 13. Martin DO, Lemke B, Birnie D, Krum H, Lee KLF, Aonuma K, Gasparini M, Starling RC, Milasinovic G, Rogers T, Sambelashvili A, Gorcsan J, Houmsse M. Investigation of a novel algorithm for synchronized left-ventricular pacing and ambulatory optimization of cardiac resynchronization therapy: Results of the adaptive CRT trial. *Heart Rhythm*. 2012;9.
 14. Ritter P, Delnoy PPH, Padeletti L, Lunati M, Naegele H, Borri-Brunetto A, Silvestre J. A randomized pilot study of optimization of cardiac resynchronization therapy in sinus rhythm patients using a peak endocardial acceleration sensor vs. standard methods. *Europace*. 2012;14:1324–1333.
 15. Gold MR, Yu Y, Singh JP, Stein KM, Birgersdöter-Green U, Meyer TE, Seth M, Ellenbogen KA. The effect of left ventricular electrical delay on AV optimization for cardiac

- resynchronization therapy. *Heart Rhythm*. 2013;10:988–993.
16. Daubert JC, Saxon L, Adamson PB, Auricchio A, Berger RD, Beshai JF, Breithard O, Brignole M, Cleland J, Delurgio DB, Dickstein K, Exner D V., Gold M, Grimm RA, Hayes DL, Israel C, Leclercq C, Linde C, Lindenfeld J, Merkely B, Mont L, Murgatroyd F, Prinzen F, Saba SF, Shinbane JS, Singh J, Tang AS, Vardas PE, Wilkoff BL, Zamorano JL. 2012 EHRA/HRS expert consensus statement on cardiac resynchronization therapy in heart failure: Implant and follow-up recommendations and management. *Heart Rhythm*. 2012;9:1524–1576.
 17. Brignole M, Auricchio A, Baron-Esquivias G, Bordachar P, Boriani G, Breithardt O-A, Cleland J, Deharo J-C, Delgado V, Elliott PM, Gorenek B, Israel CW, Leclercq C, Linde C, Mont L, Padeletti L, Sutton R, Vardas PE, ESC Committee for Practice Guidelines (CPG), Zamorano JL, Achenbach S, Baumgartner H, Bax JJ, Bueno H, Dean V, Deaton C, Erol C, Fagard R, Ferrari R, Hasdai D, Hoes AW, Kirchhof P, Knuuti J, Kolh P, Lancellotti P, Linhart A, Nihoyannopoulos P, Piepoli MF, Ponikowski P, Sirnes PA, Tamargo JL, Tendera M, Torbicki A, Wijns W, Windecker S, Document Reviewers, Kirchhof P, Blomstrom-Lundqvist C, Badano LP, Aliyev F, Bänsch D, Baumgartner H, Bsata W, Buser P, Charron P, Daubert J-C, Dobreanu D, Faerestrang S, Hasdai D, Hoes AW, Le Heuzey J-Y, Mavrakis H, McDonagh T, Merino JL, Nawar MM, Nielsen JC, Pieske B, Poposka L, Ruschitzka F, Tendera M, Van Gelder IC, Wilson CM. 2013 ESC Guidelines on cardiac pacing and cardiac resynchronization therapy: the Task Force on cardiac pacing and resynchronization therapy of the European Society of Cardiology (ESC). Developed in collaboration with the European Heart Rhythm Association. *Eur Heart J*. 2013;34:2281–329.
 18. Birnie D, Lemke B, Aonuma K, Krum H, Lee KLF, Gasparini M, Starling RC, Milasinovic G, Goresan J, Houmsse M, Abeyratne A, Sambelashvili A, Martin DO. Clinical outcomes with

synchronized left ventricular pacing: Analysis of the adaptive CRT trial. *Heart Rhythm*. 2013;10:1368–1374.

19. Filippatos G, Birnie D, Gold MR, Gerritse B, Hersi A, Jacobs S, Kusano K, Leclercq C, Mullens W, Wilkoff BL. Rationale and design of the AdaptResponse trial: a prospective randomized study of cardiac resynchronization therapy with preferential adaptive left ventricular-only pacing. *Eur J Heart Fail*. 2017;19:950–957.

Figure Legends

Figure 1: Justification for Interventricular Delay Cut-off

Panel A) Percent of CRT population with delay greater than RV-LV cutoffs, the median value was 70 msec.

Panel B) The difference in CRT response between the patients in the SmartDelay and the Fixed delay (at 120ms) cohorts by RV-LV cutoff. Cutoffs represent RV-LV durations equal to or longer than the specified cutoff value.

Data obtained from the SMART-AV Trial.




Figure 2: Study Design Flow Chart

Eligible patients (RV-LV duration ≥ 70 msec) will be randomized in the post-implant period.

Response rates (proportion of patients with $\geq 15\%$ decrease in LVESV from pre-implant echocardiogram) as well as Kansas City Cardiomyopathy Questionnaire (KCCQ), 6-Minute Hall Walk (6MW), LVESV and LVEF will be evaluated at a 6-month follow-up.




Table 1: Study inclusion criteria

Qualifying subjects are those with heart failure who CRT indications defined as those subjects who receive stable optimal pharmacologic therapy for heart failure and who meet any one of the following classifications:

- Moderate to severe heart failure (NYHA Class III-IV) with LVEF $\leq 35\%$ and QRS duration ≥ 120 msec**
- Mild heart failure (NYHA Class II) with LVEF $\leq 30\%$ and LBBB with QRS duration ≥ 130 msec**

This indication is modified in Japan to restrict the QRS cutoff to 150 msec

- Asymptomatic heart failure (NYHA Class I) with ischemic cardiomyopathy, LVEF $\leq 30\%$ and LBBB with QRS duration ≥ 130 msec**

This indication is eliminated in Japan

Subject is age 18 years old or above, or of legal age and willing and capable to give informed consent specific to each country and national laws

Subject must be indicated for a CRT-D system implant. This includes subjects who are indicated to receive an upgrade to a BSC quadripolar CRT-D device from a previously implanted device

Subject is willing and capable of complying with visits and procedures as defined by this protocol

CRT=cardiac resynchronization therapy, LBBB=left ventricular bundle branch block, LVEF=left ventricular ejection fraction

Table 2: Study exclusion criteria

Subjects with documented permanent complete AV block

Subjects with permanent or chronic AF or in AF at the time of enrollment

Subjects who have previously received cardiac resynchronization therapy with pacing in the left ventricle

Subjects on the active heart transplant list or who has or is to receive a VAD

Life expectancy shorter than 12 months due to any medical condition (e.g., cancer, uremia, liver failure)

Subject with a known or suspected sensitivity to dexamethasone acetate

Women of childbearing potential who are or plan to become pregnant during the course of the trial

Subjects currently requiring dialysis

AF=atrial fibrillation, AV=atrioventricular, **VAD=ventricular assist device**
